

Vitamin B₁₂ is bound to intrinsic factor during transit through the stomach; separation occurs in the terminal ileum in the presence of calcium, and Vitamin B₁₂ enters the mucosal cell for absorption. It is then transported by the transcobalamin binding proteins. A small amount (approximately 1% of the total amount ingested) is absorbed by simple diffusion, but this mechanism is adequate only with very large doses. Oral absorption is considered too undependable to rely on in patients with pernicious anemia or other conditions resulting in malabsorption of Vitamin B₁₂.

Cyanocobalamin is the most widely used form of Vitamin B₁₂, and has hematopoietic activity apparently identical to that of the antianemia factor in purified liver extract. Hydroxocobalamin is equally as effective as cyanocobalamin, and they share the cobalamin molecular structure.

INDICATIONS AND USAGE

Cyanocobalamin is indicated for Vitamin B₁₂ deficiencies due to malabsorption which may be associated with the following conditions:

Addisonian (pernicious) anemia

Gastrointestinal pathology, dysfunction, or surgery, including gluten enteropathy or sprue, small bowel bacterial overgrowth, total or partial gastrectomy

Fish tapeworm infestation

Malignancy of pancreas or bowel

Folic acid deficiency

It may be possible to treat the underlying disease by surgical correction of anatomic lesions leading to small bowel bacterial overgrowth, expulsion of fish tapeworm, discontinuation of drugs leading to vitamin malabsorption (see ***Drug/Laboratory Test Interactions***), use of a gluten-free diet in nontropical sprue, or administration of antibiotics in tropical sprue. Such measures remove the need for long-term administration of cyanocobalamin.

Requirements of Vitamin B₁₂ in excess of normal (due to pregnancy, thyrotoxicosis, hemolytic anemia, hemorrhage, malignancy, hepatic and renal disease) can usually be met with oral supplementation.

Cyanocobalamin injection is also suitable for the Vitamin B₁₂ absorption test (Schilling test).

CONTRAINDICATIONS

Sensitivity to cobalt and/or Vitamin B₁₂ is a contraindication.

WARNINGS

This product contains aluminum that may be toxic. Aluminum may reach toxic levels with prolonged parenteral administration if kidney function is impaired. Premature neonates are particularly at risk because their kidneys are immature, and they require large amounts of calcium and phosphate solutions, which contain aluminum.

Research indicates that patients with impaired kidney function, including premature neonates, who receive parenteral levels of aluminum at greater than 4 to 5 mcg/kg/day accumulate aluminum at levels associated with central nervous system and bone toxicity. Tissue loading may occur at even lower rates of administration.

Patients with early Leber's disease (hereditary optic nerve atrophy) who were treated with cyanocobalamin suffered severe and swift optic atrophy.

Hypokalemia and sudden death may occur in severe megaloblastic anemia which is treated intensely.

Anaphylactic shock and death have been reported after parenteral Vitamin B₁₂ administration. An

intradermal test dose is recommended before cyanocobalamin injection is administered to patients suspected of being sensitive to this drug.

This product contains benzyl alcohol. Benzyl alcohol has been reported to be associated with a fatal "Gasping Syndrome" in premature infants.

PRECAUTIONS

General

Vitamin B₁₂ deficiency that is allowed to progress for longer than three months may produce permanent degenerative lesions of the spinal cord. Doses of folic acid greater than 0.1 mg/day may result in hematologic remission in patients with Vitamin B₁₂ deficiency. Neurologic manifestations will not be prevented with folic acid, and if not treated with Vitamin B₁₂, irreversible damage will result.

Doses of cyanocobalamin exceeding 10 mcg daily may produce hematologic response in patients with folate deficiency. Indiscriminate administration may mask the true diagnosis.

Information for Patients

Patients with pernicious anemia should be instructed that they will require monthly injections of Vitamin B₁₂ for the remainder of their lives. Failure to do so will result in return of the anemia and in development of incapacitating and irreversible damage to the nerves of the spinal cord. Also, patients should be warned about the danger of taking folic acid in place of Vitamin B₁₂, because the former may prevent anemia but allow progression of subacute combined degeneration.

A vegetarian diet which contains no animal products (including milk products or eggs) does not supply any Vitamin B₁₂. Patients following such a diet should be advised to take oral Vitamin B₁₂ regularly. The need for Vitamin B₁₂ is increased by pregnancy and lactation. Deficiency has been recognized in infants of vegetarian mothers who were breast fed, even though the mothers had no symptoms of deficiency at the time.

Contains no more than 57 mcg/L of aluminum.

Laboratory Tests

During the initial treatment of patients with pernicious anemia, serum potassium must be observed closely the first 48 hours and potassium replaced if necessary.

Hematocrit, reticulocyte count, Vitamin B₁₂, folate and iron levels should be obtained prior to treatment. Hematocrit and reticulocyte counts should be repeated daily from the 5th to 7th days of therapy and then frequently until the hematocrit is normal. If folate levels are low, folic acid should also be administered. If reticulocytes have not increased after treatment or if reticulocyte counts do not continue at least twice normal as long as the hematocrit is less than 35%, diagnosis or treatment should be reevaluated. Repeat determinations of iron and folic acid may reveal a complicating illness that might inhibit the response of the marrow.

Patients with pernicious anemia have about three times the incidence of carcinoma of the stomach as the general population, so appropriate tests for this condition should be carried out when indicated.

Drug and Laboratory Test Interactions

Persons taking most antibiotics, methotrexate and pyrimethamine invalidate folic acid and Vitamin B₁₂ diagnostic blood assays.

Colchicine, para-aminosalicylic acid and heavy alcohol intake for longer than two weeks may produce malabsorption of Vitamin B₁₂.

Carcinogenesis, Mutagenesis

Long-term studies in animals to evaluate carcinogenic potential have not been done. There is no evidence from long-term use in patients with pernicious anemia that cyanocobalamin is carcinogenic. Pernicious anemia is associated with an increased incidence of carcinoma of the stomach, but this is believed to be related to the underlying pathology and not to treatment with cyanocobalamin.

Pregnancy

Adequate and well-controlled studies have not been done in pregnant women. However, Vitamin B₁₂ is an essential vitamin and requirements are increased during pregnancy. Amounts of Vitamin B₁₂ that are recommended by the Food and Nutrition Board, National Academy of Science-National Research Council for pregnant women (4 mcg daily) should be consumed during pregnancy.

Nursing Mothers

Vitamin B₁₂ is known to be excreted in human milk. Amounts of Vitamin B₁₂ that are recommended by the Food and Nutrition Board, National Academy of Science-National Research Council for lactating women (4 mcg daily) should be consumed during lactation.

Pediatric Use

Intake in children should be in the amount (0.5 to 3 mcg daily) recommended by the Food and Nutrition Board, National Academy of Science-National Research Council.

ADVERSE REACTIONS

Generalized

Anaphylactic shock and death have been reported with administration of parenteral Vitamin B₁₂ (see **WARNINGS**).

Cardiovascular

Pulmonary edema and congestive heart failure early in treatment; peripheral vascular thrombosis.

Hematological

Polycythemia vera.

Gastrointestinal

Mild transient diarrhea.

Dermatological

Itching; transitory exanthema.

Miscellaneous

Feeling of swelling of entire body.

OVERDOSAGE

No overdosage has been reported with this drug.

DOSAGE AND ADMINISTRATION

Avoid using the intravenous route. Use of this product intravenously will result in almost all of the vitamin being lost in the urine.

Pernicious Anemia

Parenteral Vitamin B₁₂ is the recommended treatment and will be required for the remainder of the

patient's life. The oral form is not dependable. A dose of 100 mcg daily for six or seven days should be administered by intramuscular or deep subcutaneous injection. If there is clinical improvement and if a reticulocyte response is observed, the same amount may be given on alternate days for seven doses, then every three to four days for another two to three weeks. By this time hematologic values should have become normal. This regimen should be followed by 100 mcg monthly for life. Folic acid should be administered concomitantly if needed.

Patients with Normal Intestinal Absorption

Where the oral route is not deemed adequate, initial treatment similar to that for patients with pernicious anemia may be indicated depending on the severity of the deficiency. Chronic treatment should be with an oral B₁₂ preparation. If other vitamin deficiencies are present, they should be treated.

Schilling Test

The flushing dose is 1,000 mcg.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

HOW SUPPLIED

NDC Number	Cyanocobalamin mcg/mL	Volume
70069- 005 -10	1,000	1 mL
70069- 172 -10		10 mL
70069- 172 -25		

1 mL multiple dose vial, packaged 25 vials per tray.

10 mL multiple dose vial, packaged as;

- 10 vials per tray and

- 25 vials per tray.

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

PROTECT FROM LIGHT

Use only if solution is clear and seal intact.

For Product Inquiry call 1-800-417-9175

Manufactured for:

Somerset Therapeutics, LLC

Hollywood, FL 33024

Made in India

Code No.: KR/DRUGS/KTK/28/289/97

ST-CYC/P/05

Revised: 09/2020

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

Cyanocobalamin Injection, USP

1 mL Container Label

NDC 70069-005-01

Contains no more than 57 mcg/L of aluminum. NDC 70069-005-01 Rx only ST-CYC15/L/07
Store at 20° to 25°C (68° to 77°F)
[see USP Controlled Room Temperature].
PROTECT FROM LIGHT
Manufactured for:
Somerset Therapeutics, LLC
Hollywood, FL 33024
Made in India
Code No.:KR/DRUGS/KTK/28/289/97 Contains Benzyl Alcohol as a Preservative

Cyanocobalamin Injection, USP
1,000 mcg/mL
For IM or SC Use
1 mL Multiple Dose Vial

No Varnish Area
14x57mm




Container Label

Cyanocobalamin Injection, USP

1 mL Carton Label

NDC 70069-005-10



The image shows a template for a 1 mL carton label for Cyanocobalamin Injection, USP. It features a large 'No Varnish Area' at the top, a central section with the product name and strength, and a bottom section with detailed information including manufacturer details, storage instructions, and a barcode.

No Varnish Area
42 x 80mm

Somerset Therapeutics LLC
For Intramuscular or Subcutaneous Use
Cyanocobalamin Injection, USP
1,000 mcg/mL
NDC 70069-005-10
25 x 1 mL Multiple Dose Vials
Rx only

Each mL contains: Cyanocobalamin 1,000 mcg; Benzyl alcohol 1.5%; Sodium chloride 0.9%; Water for Injection pH adjusted with hydrochloric acid or sodium hydroxide if necessary.
Contains no more than 57 mcg/L of Aluminum.
Not made with natural rubber latex.
Usual dosage: See package insert.
Use only if solution is clear and seal intact.
Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].
PROTECT FROM LIGHT

Manufactured for:
Somerset Therapeutics, LLC
Hollywood, FL 33024
Customer Care # 1-800-417-9175

Made in India
Code No.:KR/DRUGS/KTK/28/289/97
ST-CYC15/C25/07

Rx only
NDC 70069-005-10
25 x 1 mL Multiple Dose Vials
Cyanocobalamin Injection, USP
1,000 mcg/mL
For Intramuscular or Subcutaneous Use
CONTAINS BENZYL ALCOHOL AS A PRESERVATIVE
Somerset Therapeutics LLC




Carton Label

Cyanocobalamin Injection, USP

10 mL Container Label

NDC 70069-172-01

<p>Contains no more than 57 mcg/L of aluminum. Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. PROTECT FROM LIGHT</p> <p>Manufactured for: Somerset Therapeutics, LLC Hollywood, FL 33024</p> <p>Made in India Code No.:KR/DRUGS/KTK/28/289/97</p>	<p>NDC 70069-172-01 Rx only</p> <p>Cyanocobalamin Injection, USP</p> <p>10,000 mcg/10 mL (1,000 mcg/mL)</p> <p>For IM or SC Use</p> <p>10 mL Multiple Dose Vial</p> <p>Contains Benzyl Alcohol as a Preservative</p>	 <p>3 7 0 0 6 9 1 7 2 0 1 8</p> <p>ST-CYC12/L/04</p>	<p>No Varnish Area 24 x 7.5mm</p>
---	--	---	---------------------------------------


container Label

Cyanocobalamin Injection, USP
10 mL Carton Label – Pack of 25's
NDC 70069-172-25

<p>No Varnish Area 65 x 122mm</p> <p>Cyanocobalamin Injection, USP 10,000 mcg/10 mL (1,000 mcg/mL) For Intramuscular or Subcutaneous Use CONTAINS BENZYL ALCOHOL AS A PRESERVATIVE</p> <p>25 x 10 mL Multiple Dose Vials NDC 70069-172-25</p> <p>Rx only</p> <p>Each mL contains: Cyanocobalamin 1,000 mcg; Benzyl alcohol 1.5%; Sodium chloride 0.9%; Water for Injection q.s. pH adjusted with hydrochloric acid or sodium hydroxide if necessary. Contains no more than 57 mcg/L of Aluminum. Not made with natural rubber latex.</p> <p>Usual dosage: See package insert. Use only if solution is clear and seal intact.</p> <p>Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. PROTECT FROM LIGHT</p>	<p>Manufactured for: Somerset Therapeutics, LLC Hollywood, FL 33024 Customer Care # 1-800-417-9175</p> <p>Made in India Code No.:KR/DRUGS/KTK/28/289/97 ST-CYC12/C25/03</p>	<p>Rx only</p> <p>NDC 70069-172-25 25 x 10 mL Multiple Dose Vials</p> <p>Cyanocobalamin Injection, USP</p> <p>10,000 mcg/10 mL (1,000 mcg/mL)</p> <p>For Intramuscular or Subcutaneous Use CONTAINS BENZYL ALCOHOL AS A PRESERVATIVE</p> <p>Somerset Therapeutics LLC</p>	 <p>3 7 0 0 6 9 1 7 2 2 5 1 4</p>
--	---	--	--

carton label

Cyanocobalamin Injection, USP
10 mL Carton Label – Pack of 10's
NDC 70069-172-10



carton-label

CYANOCOBALAMIN

cyanocobalamin injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70069-005
Route of Administration	INTRAMUSCULAR, SUBCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CYANOCOBALAMIN (UNII: P6 YC3EG204) (CYANOCOBALAMIN - UNII:P6 YC3EG204)	CYANOCOBALAMIN	1000 ug in 1 mL

Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	15 mg in 1 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X)	9 mg in 1 mL
HYDROCHLORIC ACID (UNII: QTT17582CB)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	RED (Red Colored Clear Solution)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70069-005-10	25 in 1 CARTON	12/11/2015	

1	NDC:70069-005-01	1 mL in 1 VIAL; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA206503	12/11/2015		

CYANOCOBALAMIN
cyanocobalamin injection

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70069-172
Route of Administration	INTRAMUSCULAR, SUBCUTANEOUS		

Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	CYANOCOBALAMIN (UNII: P6YC3EG204) (CYANOCOBALAMIN - UNII:P6YC3EG204)	CYANOCOBALAMIN	1000 ug in 1 mL

Inactive Ingredients		
	Ingredient Name	Strength
	BENZYL ALCOHOL (UNII: LKG8494WBH)	15 mg in 1 mL
	SODIUM CHLORIDE (UNII: 451W47IQ8X)	9 mg in 1 mL
	HYDROCHLORIC ACID (UNII: QTT17582CB)	
	SODIUM HYDROXIDE (UNII: 55X04QC32I)	
	WATER (UNII: 059QF0KO0R)	

Product Characteristics			
Color	RED (Red Colored Clear Solution)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70069-172-10	10 in 1 CARTON	06/27/2017	
1	NDC:70069-172-01	10 mL in 1 VIAL; Type 0: Not a Combination Product		
2	NDC:70069-172-25	25 in 1 CARTON	06/27/2017	
2	NDC:70069-172-01	10 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206503	06/27/2017	

Labeler - Somerset Therapeutics, LLC (079947873)

Registrant - Somerset Therapeutics, LLC (079947873)

Establishment

Name	Address	ID/FEI	Business Operations
Wintac Limited		677236695	ANALYSIS(70069-005, 70069-172) , LABEL(70069-005, 70069-172) , MANUFACTURE(70069-005, 70069-172) , PACK(70069-005, 70069-172)

Revised: 9/2020

Somerset Therapeutics, LLC